

# Louth: Antimicrobial Guidelines - Louth Hospitals: Antimicrobial Guidelines: Voriconazole Prescribing Aid

## Voriconazole Prescribing Aid

Please see BNF or SPC for full prescribing information.

### Dose :

**IV:** Loading dose 6 mg/kg IV every 12 hours for 2 doses, then maintenance dose 4 mg/kg IV every 12 hours. <sup>1-3</sup>

**Oral:** **N.B.** Oral voriconazole **must** be taken on an empty stomach for effect.

If patient is on NG feed, a 3 hour feedbreak is required for each dose: Stop NG feed 2 hours before dose, restart NG feed 1 hour after dose. <sup>4</sup>

**Adult (body-weight up to 40 kg):** Initially 200 mg PO every 12 hours for 2 doses, then 100 mg PO every 12 hours, increased if necessary to 150 mg PO every 12 hours. <sup>1-3</sup>

**Adult (body-weight 40 kg and above):** Initially 400 mg PO every 12 hours for 2 doses, then 200 mg PO every 12 hours, increased if necessary to 300 mg PO every 12 hours. <sup>1-3</sup>

### Dose in Renal Impairment :

No dose adjustment required in renal impairment or CRRT. <sup>5</sup> When CrCl < 50 mL/min, accumulation of the intravenous vehicle may occur. Monitor renal function if patient on IV treatment. <sup>1,5</sup>

Use oral route unless risk/benefit justifies use of IV, e.g. check if patient absorbing PO/NG medicines.

### Drug Interactions :

Voriconazole can increase QTc – monitor, particularly if patient on other medication which can increase QTc. <sup>6</sup>

Simvastatin: Voriconazole increases exposure to simvastatin. Suggest HOLD (combination contra-indicated). <sup>6</sup>

Atorvastatin: Voriconazole increases exposure to atorvastatin. Suggest HOLD (manufacturer advises to avoid or use a lower maximum dose and monitor for rhabdomyolysis). <sup>6</sup>

Benzodiazepines: Voriconazole is likely to increase the concentration of benzodiazepines and lead to a prolonged sedative effect, dose reduction may be required. <sup>6</sup>

Phenytoin decreases the exposure to voriconazole and voriconazole increases exposure to phenytoin. Manufacturer advises avoid or adjust voriconazole dose and monitor phenytoin level. <sup>6</sup>

Multiple other interactions, see BNF or SPC.

### Therapeutic Drug Monitoring :

Check trough level after 3 to 5 days. <sup>7</sup>

Use serum bottle brown top tube

Repeat level in the second week to ensure it is in therapeutic range. <sup>7</sup>

In practice, send level on Monday so that Laboratory Referrals can send out on Tuesday morning for processing on Wednesday. Result usually available by Friday evening.

### Other Monitoring :

Monitor ALT and AST <sup>1-3</sup>

Monitor for rash – risk of SJS or TEN with voriconazole. <sup>2,3</sup>

Reports of prolonged visual adverse reactions, including blurred vision, optic neuritis and papilloedema. <sup>2,3</sup>

Risk of phototoxicity.<sup>1-3</sup>

**References :**

British National Formulary. Available from [www.medicinescomplete.com](http://www.medicinescomplete.com) , accessed 22/2/21.

Fresenius Kabi. Summary of Product Characteristics for Voriconazole 200mg powder for solution for infusion. 2020. Available from [www.hpra.ie](http://www.hpra.ie) , accessed 22/2/21.

Pfizer. Summary of Product Characteristics for Vfend® 50mg and 200mg tablets. 2020. Available from [www.hpra.ie](http://www.hpra.ie) , accessed 22/2/21.

The NEWT Guidelines. Available from [www.newtguidelines.com](http://www.newtguidelines.com) , accessed 22/2/21.

The Renal Drug Database. Available from [www.renaldrugdatabase.com](http://www.renaldrugdatabase.com) , accessed 22/2/21.

Stockley's Drug Interactions. Available from [www.medicinescomplete.com](http://www.medicinescomplete.com) , accessed 22/2/21.

ECIL 6 Meeting. Triazole Antifungal Therapeutic Drug Monitoring. Final slide set posted on ECIL website Dec 8<sup>th</sup> , 2015. On file in OLOL Pharmacy Department.

Document prepared by: Carmel McKenna; Checked by: Catriona Campbell; OLOL Pharmacy Department Feb 2021.